

Anti-CAT Polyclonal Antibody

Summary

Catalog No. PHC05301

Host species Rabbit

Tested applications ELISA: 1:4000-1:8000, IHC: 1:50-1:100, WB: 1:1000-1:4000

Species reactivity Human, Mouse, Rat

Immunogen E. coli - derived recombinant Human CAT (Met1-Leu527).

Form Liquid

Storage buffer 0.01M PBS, pH 7.4, 50% Glycerol, 0.05% Proclin 300.

Clonality Polyclonal

Isotype IgG

Applications ELISA, IHC, WB

Target Catalase,CAT

Purification Purified by antigen affinity column.

Accession P04040

Use a manual defrost freezer and avoid repeated freeze thaw cycles. Store


Stability and Storage at 2 to 8°C for frequent use. Store at -20 to -80°C for twelve months from

the date of receipt.


Note For research use only.

Data Image

Recombinant Proteins & Antibodies



Western blot



Immunocytochemistry/ Immunofluorescence

Various lysates were subjected to SDS PAGE followed by western blot with CAT antibody (PHC05301) at 0.5µg/ml.

Lane 1: 293T cell lysate Lane 2: A549 cell lysate Lane 3: HepG2 cell lysate Lane 4: A2780 cell lysate

Second Ab: Goat Anti-Rabbit IgG H&L Polyclonal antibody, HRP (PTB96431) at 0.1 µg/mL.


Predict MW: 60 kDa

Observed MW: 60,62 kDa

CAT in HepG2 Cell Line.

The HepG2 cells were fixed with 4% paraformaldehyde (20 min), and then blocked with 5% goat serum for 1h. And the cells were incubated for 2h at 37°C with CAT (PHC05301) at 10 µg/ml. The section was then incubated with Goat Anti-Rabbit IgG (Alexa Fluor-488) preabsorbed at 1/100 dilution (Shown in green) for 1 hour at room temperature. Nuclear DNA was labelled with DAPI (shown in blue).

Recombinant Proteins & Antibodies



Western Blot

Recombinant Protein lysates were subjected to SDS PAGE followed by western blot with CAT antibody (PHC05301) at 1 μg/ml.

Lane 1: Recombinant Protein

Second Ab: Goat Anti-Rabbit IgG H&L Polyclonal antibody, HRP (PTB96431) at 0.1 µg/mL.

Predict MW: 62 kDa Observed MW: 62 kDa