

# Anti-Human TBCA Polyclonal Antibody


## Summary

---

<b>Catalog No.</b>	PHB15901
<b>Host species</b>	Rabbit
<b>Tested applications</b>	ELISA: 1:4000-1:8000, IHC: 1:50-1:100, WB: 1:1000-1:4000
<b>Species reactivity</b>	Human
<b>Immunogen</b>	E. coli - derived recombinant Human TBCA (Ala2-Ala108).
<b>Form</b>	Liquid
<b>Storage buffer</b>	0.01M PBS, pH 7.4, 50% Glycerol, 0.05% Proclin 300.
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Applications</b>	ELISA, IHC, WB
<b>Target</b>	Tubulin-specific chaperone A, TBCA, Tubulin-folding cofactor A, CFA, TCP1-chaperonin cofactor A
<b>Purification</b>	Purified by antigen affinity column.
<b>Accession</b>	O75347
<b>Stability and Storage</b>	Use a manual defrost freezer and avoid repeated freeze thaw cycles. Store at 2 to 8°C for frequent use. Store at -20 to -80°C for twelve months from the date of receipt.
<b>Note</b>	For research use only.

## Data Image

---



Western Blot

Various lysates were subjected to SDS PAGE followed by western blot with TBICA antibody (PHB15901) at 1  $\mu$ g/ml.

Lane 1: Hela cell lysate

Lane 2: 293T cell lysate

Lane 3: Jurkat cell lysate

Lane 4: Lncap cell lysate


Lane 5: HepG2 cell lysate

Lane 6: A431 cell lysate

Second Ab: Goat Anti-Rabbit IgG H&L Polyclonal antibody, HRP (PTB96431) at 0.1  $\mu$ g/mL.

Predict MW: 16 kDa

Observed MW: 16 kDa



Western Blot

Recombinant Protein lysates were subjected to SDS PAGE followed by western blot with TBICA antibody (PHB15901) at 1  $\mu$ g/ml.

Lane 1: Recombinant protein

Second Ab: Goat Anti-Rabbit IgG H&L Polyclonal antibody, HRP (PTB96431) at 0.1  $\mu$ g/mL.

Predict MW: 25 kDa

Observed MW: 25 kDa