

## Anti-SUMO2 Polyclonal Antibody

## Summary

Catalog No. PHF46901

**Host species** Rabbit

**Tested applications** ELISA: 1:4000-1:8000, IHC: 1:50-1:100, WB: 1:1000-1:4000

**Species reactivity** Human, Mouse, Rat

**Immunogen** E. coli - derived recombinant Human SUMO2 (Met1-Tyr95).

**Form** Liquid

Storage buffer 0.01M PBS, pH 7.4, 50% Glycerol, 0.05% Proclin 300.

**Clonality** Polyclonal

**Isotype** IgG

**Applications** ELISA, IHC, WB

Ubiquitin-like protein SMT3B,SMT3B,SUMO-3,Smt3B,Sentrin-2,SMT3

**Target** homolog 2,SMT3H2,SUMO2,HSMT3,Small ubiquitin-related modifier

2,SUMO-2

**Purification** Purified by antigen affinity column.

Accession P61956

Use a manual defrost freezer and avoid repeated freeze thaw cycles. Store


Stability and Storage at 2 to 8°C for frequent use. Store at -20 to -80°C for twelve months from

the date of receipt.


Note For research use only.

## Data Image

## Recombinant Proteins & Antibodies



WESTERN BLOT



Immunocytochemistry/ Immunofluorescence

Recombinant Protein lysates were subjected to SDS PAGE followed by western blot with SUMO2 antibody (PHF46901) at 1 µg/ml.

Lane 1: Recombinant protein

Second Ab: Goat Anti-Rabbit IgG H&L Polyclonal antibody, HRP (PTB96431) at 0.1 µg/mL.

Predict MW: 13 kDa Observed MW: 13 kDa

SUMO2 in 293T Cell Line.

The 293T cells were fixed with 4% paraformaldehyde (20 min), and then blocked with 5% goat serum for 1h. And the cells were incubated for 2h at 37°C with SUMO2 (PHF46901) at 4.8 µg/ml. The section was then incubated with Goat Anti-Rabbit IgG (Alexa Fluor-488) preabsorbed at 1/100 dilution (Shown in green) for 1 hour at room temperature. Nuclear DNA was labelled with DAPI (shown in blue).