

Anti-ETFA Antibody (R3D47)


Summary

Catalog No.	RHD01401
Clone ID	R3D47
Host species	Mouse
Tested applications	ELISA: 1:10000, FCM: 1:200-1:400, IHC: 1:200-1:1000, WB: 1:500-1:2000
Species reactivity	Human
Form	Liquid
Storage buffer	0.01M PBS, pH 7.4, 0.05% Sodium Azide.
Concentration	1 mg/ml
Purity	>95% as determined by SDS-PAGE.
Clonality	Monoclonal
Isotype	IgG1
Applications	ELISA, FCM, IHC, WB
Target	Alpha-ETF, ETFA, Electron transfer flavoprotein subunit alpha, mitochondrial
Purification	Protein A/G purified from cell culture supernatant.
Endotoxin level	Please contact with the lab for this information.
Accession	P13804
Stability and Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. Store at 4°C short term (1-2 weeks). Store at -20°C 12 months. Store at -80°C long term.

Note


For research use only.

Data Image




Flow Cytometry

Flow cytometric analysis of HeLa cells using ETFA mouse mAb (green) and negative control (red).




Flow Cytometry

Flow cytometric analysis of HepG2 cells using ETFA mouse mAb (green) and negative control (red).




Immunohistochemical

Immunohistochemical analysis of paraffin-embedded human cervical carcinoma tissues using ETFA mouse mAb with DAB staining.




Immunohistochemical

Immunohistochemical analysis of paraffin-embedded human bladder cancer tissues using ETFA mouse mAb with DAB staining.




Immunohistochemical

Immunohistochemical analysis of paraffin-embedded human breast cancer tissues using ETFA mouse mAb with DAB staining.




Immunohistochemical

Immunohistochemical analysis of paraffin-embedded human breast cancer tissues using ETFA mouse mAb with DAB staining.



Immunohistochemical

Immunohistochemical analysis of paraffin-embedded human rectal cancer tissues using ETFA mouse mAb with DAB staining.



Western blot

Western blot analysis using ETFA mouse mAb against .HepG2 (1), A431 (2),Hek293 (3),Hela (4) and MCF-7 (5) cell lysate.