

# Anti-NCF2 Antibody (R3F46)

## Summary

---

|                            |  |
|----------------------------|--|
| <b>Catalog No.</b>         | RHD36202   |
| <b>Clone ID</b>            | R3F46  |
| <b>Host species</b>        | Mouse  |
| <b>Tested applications</b> | IHC: 1:100-1:200   |
| <b>Species reactivity</b>  | Human  |
| <b>Form</b>                | Liquid   |
| <b>Storage buffer</b>      | 0.01M PBS, pH 7.4, 0.05% Sodium Azide.   |
| <b>Concentration</b>       | 1 mg/ml  |
| <b>Purity</b>              | >95% as determined by SDS-PAGE.  |
| <b>Clonality</b>           | Monoclonal   |
| <b>Isotype</b>             | IgG1   |
| <b>Applications</b>        | IHC  |
| <b>Target</b>              | Neutrophil NADPH oxidase factor 2, NOXA2, p67-phox, NADPH oxidase activator 2, 67 kDa neutrophil oxidase factor, Neutrophil cytosol factor 2, NCF-2, NCF2, P67PHOX |
| <b>Purification</b>        | Protein A/G purified from cell culture supernatant.  |
| <b>Endotoxin level</b>     | Please contact with the lab for this information.  |
| <b>Accession</b>           | P19878   |

**Stability and Storage**

Use a manual defrost freezer and avoid repeated freeze-thaw cycles.


Store at 4°C short term (1-2 weeks). Store at -20°C 12 months. Store at -80°C long term.

**Note**

For research use only.

## Data Image

---



Immunohistochemical

IHC-P analysis of lung tissue by NCF2 antibody (RHD36202). IHC-P was performed using sections of the formalin-fixed paraffin-embedded lung tissue; Result: Macrophages are positively stained at the cytoplasm.