


# Anti-CDA Antibody (R3J38)

## Summary

---


<b>Catalog No.</b>	RHE00202
<b>Clone ID</b>	R3J38
<b>Host species</b>	Mouse
<b>Tested applications</b>	IHC: 1:50-1:250, IP: 1:100-1:200, WB: 1:500-1:1000
<b>Species reactivity</b>	Human
<b>Form</b>	Liquid
<b>Storage buffer</b>	0.01M PBS, pH 7.4, 0.05% Sodium Azide.
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	>95% as determined by SDS-PAGE.
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	IgG1
<b>Applications</b>	IHC, IP, WB
<b>Target</b>	Cytidine deaminase, CDD, Cytidine aminohydrolase, CDA
<b>Purification</b>	Protein A/G purified from cell culture supernatant.
<b>Endotoxin level</b>	Please contact with the lab for this information.
<b>Accession</b>	P32320
<b>Stability and Storage</b>	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. Store at 4°C short term (1-2 weeks). Store at -20°C 12 months. Store at -80°C long term.
<b>Note</b>	For research use only.

## Data Image




Immunohistochemical

IHC-P analysis of spleen tissue by CDA antibody (RHE00202). IHC-P was performed using sections of the formalin-fixed paraffin-embedded spleen tissue; Result: Cells in red pulp are positively stained.



Immunoprecipitation

Lane 1: SK-OV-3 lysate; Lane 2: CDA immunoprecipitated from SK-OV-3 lysate by RHE00202; Lane 3: The same as Lane 2 but use IgG isotype control antibody; Result: RHE00202 can immunoprecipitate CDA;



Western blot

Western blot analysis using CDA mouse mAb against HeLa lysate and 30 µg SK-OV-3 lysate