Anti-CD68 Antibody (R3J56)

Summary

Catalog No. RHE04901

Clone ID R3J56

Host species Mouse

Tested applications IF: 1:50-1:200, IHC: 1:50-1:100

Species reactivity Human, Mouse

Form Liquid

Storage buffer 0.01M PBS, pH 7.4, 0.5% BSA, 0.05% Sodium Azide and 50% Glycerol.

Concentration 1 mg/ml

Purity >95% as determined by SDS-PAGE.

Clonality Monoclonal

Isotype IqG1

Applications IF, IHC

Target CD68, Gp110, Macrosialin

Purification Protein A/G purified from cell culture supernatant.

Endotoxin level Please contact with the lab for this information.

Accession P34810

Use a manual defrost freezer and avoid repeated freeze-thaw cycles.


Stability and Storage Store at 4°C short term (1-2 weeks). Store at -20°C 12 months. Store at -

80°C long term.

Note For research use only.




Data Image




Immunofluorescence analysis of CD68 in Human lungcancer tissue using CD68 antibody(red)and DAPI (blue).

Immunofluorescence



Immunofluorescence analysis of CD68 in mouse spleen tissue using CD68 antibody(6F3)(red),and DAPI (blue).

Immunofluorescence




Immunohistochemical


Immunohistochemistry analysis of paraffin-embedded Human Tonsil tissue using CD68 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval. Negative control was used by secondary antibody only.




Recombinant Proteins & Antibodies



Immunohistochemical



Immunohistochemical



Immunohistochemical

Immunohistochemical analysis of paraffin-embedded Human tonsils using CD68 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.Negative control was used by secondary antibody only.

Immunohistochemistry analysis of paraffin-embedded Human tonsil tissue using CD68 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Immunohistochemistry analysis of paraffin-embedded Human tonsil tissue using CD68 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.