

# Anti-AGR3 Antibody (R3Z48)

## Summary

---

<b>Catalog No.</b>	RHK87402
<b>Clone ID</b>	R3Z48
<b>Host species</b>	Mouse
<b>Tested applications</b>	IHC: 1:100-1:200, IP: 1:100-1:200
<b>Species reactivity</b>	Human
<b>Form</b>	Liquid
<b>Storage buffer</b>	0.01M PBS, pH 7.4, 0.05% Sodium Azide.
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	>95% as determined by SDS-PAGE.
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	IgG1
<b>Applications</b>	IHC, IP
<b>Target</b>	AGR3, AG3, PDIA18, Protein disulfide isomerase family A, member 18, Breast cancer membrane protein 11, AG-3, Anterior gradient protein 3, hAG-3, BCMP11, Anterior gradient 3 homolog
<b>Purification</b>	Protein A/G purified from cell culture supernatant.
<b>Endotoxin level</b>	Please contact with the lab for this information.
<b>Accession</b>	Q8TD06

**Stability and Storage**

Use a manual defrost freezer and avoid repeated freeze-thaw cycles.


Store at 4°C short term (1-2 weeks). Store at -20°C 12 months. Store at -80°C long term.

**Note**

For research use only.


## Data Image

---




Immunohistochemical

IHC-P analysis of gall bladder tissue by AGR3 antibody (RHK87402). IHC-P was performed using sections of the formalin-fixed paraffin-embedded gall bladder tissue; Result: Glandular cells are positively stained at the cytoplasm and cell membrane.



Immunohistochemical

IHC-P analysis of fallopian tube tissue by AGR3 antibody (RHK87402). IHC-P was performed using sections of the formalin-fixed paraffin-embedded fallopian tube tissue; Result: Ciliated cells are positively stained at the cytoplasm and cell membrane.



Lane 1: T-47D lysate ; Lane 2: AGR3  
immunoprecipitated from T-47D lysate by  
RHK87402; Lane 3: The same Lane 2 but use IgG  
isotype control antibody ; Result: RHK87402 can  
immunoprecipitate AGR3;