

Anti-Human EGFR/ERBB1/HER1 Antibody (11F8)

Summary

Catalog No. FHB86910

Clone ID 11F8

Host species Human

Conjugation Unconjugated

Species reactivity Human

Form Liquid

Storage buffer 0.01M PBS, pH 7.4.

Concentration 0.8 mg/ml

Purity >95% as determined by SDS-PAGE.

Clonality Monoclonal

Isotype IgG1, kappa

Applications ELISA, FCM, WB

Target HER1, Receptor tyrosine-protein kinase erbB-1, Proto-oncogene c-ErbB-1,

Epidermal growth factor receptor, EGFR, ERBB1, ERBB

Purification Protein A/G purified from cell culture supernatant.

Endotoxin level Please contact with the lab for this information.

Accession P00533

Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Stability and Storage Store at 4°C short term (1-2 weeks). Store at -20°C 12 months. Store at -


80°C long term.




Note

For research use only.

Data Image



Flow-cytometry




Flow-cytometry

Flow-cytometry using anti-human EGFR antibody. HeLa cells were stained with an irrelevant antibody (Blue Histogram) or an anti-human EGFR antibody monoclonal antibody (Catalog # FHB86910 ,Green Histogram) at a concentration of 5 μg/ml for 30 mins at RT. After washing, bound antibody was detected using a FITC conjugated goat anti-human antibody (Catalog # PHB96441) and cells analysed on a NovoCyte Flow Cytometer.

Flow-cytometry using anti-human EGFR antibody.A431 cells were stained with an irrelevant antibody (Blue Histogram) or an anti-human EGFR antibody monoclonal antibody (Catalog # FHB86910 ,Green Histogram) at a concentration of 5 μg/ml for 30 mins at RT. After washing, bound antibody was detected using a FITC conjugated goat anti-human antibody (Catalog # PHB96441) and cells analysed on a NovoCyte Flow Cytometer.

Recombinant Proteins & Antibodies



Flow-cytometry

Flow-cytometry using anti-human EGFR antibody.SK-BR-3 cells were stained with an irrelevant antibody (Blue Histogram) or an antihuman EGFR antibody monoclonal antibody (Catalog # FHB86910, Green Histogram) at a concentration of 5 µg/ml for 30 mins at RT. After washing, bound antibody was detected using a FITC conjugated goat anti-human antibody (Catalog # PHB96441) and cells analysed on a NovoCyte Flow Cytometer.