

# Anti-Human CD136/MST1R Antibody (RON8)

## Summary


---

<b>Catalog No.</b>	FHG03410
<b>Clone ID</b>	RON8
<b>Host species</b>	Mouse
<b>Conjugation</b>	Unconjugated
<b>Species reactivity</b>	Human
<b>Form</b>	Liquid
<b>Storage buffer</b>	0.01M PBS, pH 7.4.
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	>95% as determined by SDS-PAGE.
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	IgG1, kappa
<b>Applications</b>	ELISA, FCM, WB
<b>Target</b>	CD136, Macrophage-stimulating protein receptor, PTK8, MST1R, p185-Ron, RON, CDw136, MSP receptor, Protein-tyrosine kinase 8
<b>Purification</b>	Protein A/G purified from cell culture supernatant.
<b>Endotoxin level</b>	Please contact with the lab for this information.
<b>Accession</b>	Q04912
<b>Stability and Storage</b>	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. Store at 4°C short term (1-2 weeks). Store at -20°C 12 months. Store at -80°C long term.

**Note**


For research use only.

## Data Image



Flow-cytometry

Flow-cytometry using anti-human CD136 antibody. HaCaT cells were stained with an irrelevant antibody (Blue Histogram) or an anti-human CD136 antibody monoclonal antibody (Catalog # FHG03410, Green Histogram) at a concentration of 5 µg/ml for 30 mins at RT. After washing, bound antibody was detected using a FITC conjugated goat anti-mouse antibody (Catalog # PMB96441) and cells analysed on a NovoCyt Flow Cytometer.



Flow-cytometry

Flow-cytometry using anti-human CD136 antibody. SW480 cells were stained with an irrelevant antibody (Blue Histogram) or an anti-human CD136 antibody monoclonal antibody (Catalog # FHG03410, Green Histogram) at a concentration of 5 µg/ml for 30 mins at RT. After washing, bound antibody was detected using a FITC conjugated goat anti-mouse antibody (Catalog # PMB96441) and cells analysed on a NovoCyt Flow Cytometer.