

Anti-Human CD184/CXCR4 Antibody (SAA0068)

Summary

Catalog No. FHF39920

Clone ID SAA0068

Host species Mouse

Conjugation Unconjugated

Species reactivity Human

Form Liquid

Storage buffer 0.01M PBS, pH 7.4.

Concentration 1 mg/ml

Purity >95% as determined by SDS-PAGE.

Clonality Monoclonal

Isotype IgG1

Applications FCM

Stromal cell-derived factor 1 receptor, LCR1, HM89, LPS-associated

protein 3, LAP-3, CXC-R4, CXCR-4, SDF-1 receptor, CD184, NPYRL, LESTR,

Target Fusin, FB22, C-X-C chemokine receptor type 4, CXCR4,

Lipopolysaccharide-associated protein 3, Leukocyte-derived seven

transmembrane domain receptor

Purification Protein A/G purified from cell culture supernatant.

Endotoxin level Please contact with the lab for this information.

Accession P61073



Recombinant Proteins & Antibodies

Use a manual defrost freezer and avoid repeated freeze-thaw cycles.


Store at 4°C short term (1-2 weeks). Store at -20°C 12 months. Store at -

80°C long term.


Note For research use only.

Data Image

Stability and Storage



Flow-cytometry




Flow-cytometry

Flow-cytometry using anti-human CD184 antibody. HeLa cells were stained with an irrelevant antibody (Blue Histogram) or an anti-human CD184 antibody monoclonal antibody (Catalog # FHF39920, Green Histogram) at a concentration of 5 µg/ml for 30 mins at RT. After washing, bound antibody was detected using a FITC conjugated goat anti-mouse antibody (Catalog # PMB96441) and cells analysed on a NovoCyte Flow Cytometer.

Flow-cytometry using anti-human CD184 antibody. Jurkat cells were stained with an irrelevant antibody (Blue Histogram) or an anti-human CD184 antibody monoclonal antibody (Catalog # FHF39920, Green Histogram) at a concentration of 5 µg/ml for 30 mins at RT. After washing, bound antibody was detected using a FITC conjugated goat anti-mouse antibody (Catalog # PMB96441) and cells analysed on a NovoCyte Flow Cytometer.

Recombinant Proteins & Antibodies



Flow-cytometry

Flow-cytometry using anti-human CD184 antibody. Human peripheral blood lymphocytes were stained with an irrelevant antibody (Blue Histogram) or an anti-human CD184 antibody monoclonal antibody (Catalog # FHF39920, Green Histogram) at a concentration of 5 µg/ml for 30 mins at RT. After washing, bound antibody was detected using a FITC conjugated goat anti-mouse antibody (Catalog # PMB96441) and cells analysed on a NovoCyte Flow Cytometer.