

Anti-Human CD340/ERBB2/HER2 Antibody (MAB 2C4)

Summary

Catalog No. FHC09620

Clone ID MAB 2C4

Host species Mouse

Conjugation Unconjugated

Species reactivity Human

Form Liquid

Storage buffer 0.01M PBS, pH 7.4.

Concentration 1 mg/ml

Purity >95% as determined by SDS-PAGE.

Clonality Monoclonal

Isotype IgG2a, kappa

Applications FCM

p185erbB2, HER2, NGL, Tyrosine kinase-type cell surface receptor HER2,

NEU, MLN 19, Proto-oncogene Neu, MLN19, ERBB2, Proto-oncogene c-

ErbB-2, CD340, Receptor tyrosine-protein kinase erbB-2, Metastatic lymph

node gene 19 protein

Purification Protein A/G purified from cell culture supernatant.

Endotoxin level Please contact with the lab for this information.

Accession P04626

Target



Recombinant Proteins & Antibodies


Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Stability and Storage Store at 4°C short term (1-2 weeks). Store at -20°C 12 months. Store at -


80°C long term.

Note For research use only.

Data Image



Flow-cytometry




Flow-cytometry


Flow-cytometry using anti-human CD340 antibody.MCF7 cells were stained with an irrelevant antibody (Blue Histogram) or an anti-human CD340 antibody monoclonal antibody (Catalog # FHC09620, Green Histogram) at a concentration of 5 µg/ml for 30 mins at RT. After washing, bound antibody was detected using a FITC conjugated goat anti-mouse antibody (Catalog # PMB96441) and cells analysed on a NovoCyte Flow Cytometer.

Flow-cytometry using anti-human CD340 antibody.MDA-MB-231 cells were stained with an irrelevant antibody (Blue Histogram) or an anti-human CD340 antibody monoclonal antibody (Catalog # FHC09620, Green Histogram) at a concentration of 5 µg/ml for 30 mins at RT. After washing, bound antibody was detected using a FITC conjugated goat anti-mouse antibody (Catalog # PMB96441) and cells analysed on a NovoCyte Flow Cytometer.


Recombinant Proteins & Antibodies



Flow-cytometry



Flow-cytometry



Flow-cytometry using anti-human CD340 antibody.MDA-MB-231 cells were stained with an irrelevant antibody (Blue Histogram) or an anti-human CD340 antibody monoclonal antibody (Catalog # FHC09620 ,Green Histogram) at a concentration of 5 μ g/ml for 30 mins at RT. After washing, bound antibody was detected using a FITC conjugated goat anti-mouse antibody (Catalog # PMB96441) and cells analysed on a NovoCyte Flow Cytometer.

Flow-cytometry using anti-human CD340 antibody.SK-BR-3 cells were stained with an irrelevant antibody (Blue Histogram) or an anti-human CD340 antibody monoclonal antibody (Catalog # FHC09620 ,Green Histogram) at a concentration of 5 µg/ml for 30 mins at RT. After washing, bound antibody was detected using a FITC conjugated goat anti-mouse antibody (Catalog # PMB96441) and cells analysed on a NovoCyte Flow Cytometer.

The purity of this product is >95% as determined by SEC-HPLC.