

Anti-Human CD49c/ITGA3 Antibody (84H7)

Summary

Catalog No. FHD69110

Clone ID 84H7

Host species Mouse

Conjugation Unconjugated

Species reactivity Human

Form Liquid

Storage buffer 0.01M PBS, pH 7.4.

Concentration 1 mg/ml

Purity >95% as determined by SDS-PAGE.

Clonality Monoclonal

Isotype IgG1, kappa

Applications ELISA, FCM, IP, WB

Target ITGA3, CD49 antigen-like family member C, VLA-3 subunit alpha, GAPB3,

CD49c, FRP-2, Integrin alpha-3, MSK18, Galactoprotein B3

Purification Protein A/G purified from cell culture supernatant.

Endotoxin level Please contact with the lab for this information.

Accession P26006

Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Stability and Storage Store at 4°C short term (1-2 weeks). Store at -20°C 12 months. Store at -


80°C long term.




Note

For research use only.

Data Image



Flow-cytometry




Flow-cytometry

Flow-cytometry using anti-human CD49c antibody. HeLa cells were stained with an irrelevant antibody (Blue Histogram) or an anti-human CD49c antibody monoclonal antibody (Catalog # FHD69110, Green Histogram) at a concentration of 5 µg/ml for 30 mins at RT. After washing, bound antibody was detected using a FITC conjugated goat anti-mouse antibody (Catalog # PMB96441) and cells analysed on a NovoCyte Flow Cytometer.

Flow-cytometry using anti-human CD49c antibody.KG-1 cells were stained with an irrelevant antibody (Blue Histogram) or an anti-human CD49c antibody monoclonal antibody (Catalog # FHD69110, Green Histogram) at a concentration of 5 µg/ml for 30 mins at RT. After washing, bound antibody was detected using a FITC conjugated goat anti-mouse antibody (Catalog # PMB96441) and cells analysed on a NovoCyte Flow Cytometer.

Recombinant Proteins & Antibodies



Flow-cytometry

Flow-cytometry using anti-human CD49c antibody.MG-63 cells were stained with an irrelevant antibody (Blue Histogram) or an antihuman CD49c antibody monoclonal antibody (Catalog # FHD69110, Green Histogram) at a concentration of 5 µg/ml for 30 mins at RT. After washing, bound antibody was detected using a FITC conjugated goat anti-mouse antibody (Catalog # PMB96441) and cells analysed on a NovoCyte Flow Cytometer.